

SUPPLEMENTARY MATERIAL

Importance of Angiogenin and Endothelial Progenitor Cells after Rehabilitation both in Ischemic Stroke Patients and in a Mouse Model of Cerebral Ischemia.

Marina Gabriel-Salazar^a, Anna Morancho^a, Susana Rodriguez^b, Xavi Buxó^b, Nicolás García^b, Guillem Colell^a, Albert Fernandez^a, Dolors Giralt^a, Alejandro Bustamante^a, Joan Montaner^a, Anna Rosell^{a*}.

^aNeurovascular Research Laboratory and Neurology Department, Vall d'Hebron Research Institute, Universitat Autònoma de Barcelona, Barcelona, Spain


^bUnidad de Rehabilitación Neurológica y Daño Cerebral, Hospital Vall d'Hebron, Barcelona, Spain.

Supplementary Table 1: Demographic and clinical data from postmortem samples of ischemic stroke patients.

Sex	Age	T-O-D (h)	PMI (h)	IHC	WB
Female	83	100	14.5		X
Male	84	40	8	X	X
Female	73	44	4	X	
Male	75	19	5	X	X

Abbreviations: T-O-D: time from onset of stroke symptoms to death; h:hours; PMI: postmortem interval (from death to brain samples collection) ;IHC: immunohistochemistry; WB: western blot.

Supplementary Figure 1: Full image of western blot shown in Fig.3A.



Supplementary Figure 2: Full image of western blot shown in Fig.6A.

