

## Supplementary Material

### ***BrLAS*, a *GRAS* Transcription Factor from *Brassica rapa*, is Involved in Drought Stress Tolerance in Transgenic *Arabidopsis***

Pan Li, Bin Zhang, Tongbing Su, Peirong Li, Xiaoyun Xin, Weihong Wang, Xiuyun Zhao; Yangjun Yu, Deshuang Zhang, Shuancang Yu\*, and Fenglan Zhang\*

\*Correspondence: Shuancang Yu, yushuancang@nercv.org


Fenglan Zhang, zhangfenglan@nercv.org

Supplementary Table S1. Primers used in this study.


Supplementary Table S2. Comparisons of the phenotypes of wild-type and *BrLAS*-overexpressing plants.

Supplementary Figure S1 The expression profile of *BrLAS* under normal conditions in *B.rapa*.

Supplementary Figure S2. Lateral root growth in 35S: *BrLAS* transgenic compared to wild-type (Col-0) plants.



Supplementary Figure S1 The expression profile of *BrLAS* under normal conditions in *B.rapa*. The expression patterns of *BrLAS* under normal conditions using solution without ABA, NaCl, PEG, or H<sub>2</sub>O<sub>2</sub>. Expression data of the 0h sample was normalized to 1



**Supplementary Figure S2 Lateral root growth in 35S: *BrLAS* transgenic compared to wild-type (Col-0) plants.** A relatively larger number of lateral roots were observed in the transgenic compared to wild-type plants.