

Supplementary Material

Timing of sandeel spawning and hatching off the east coast of Scotland

Alan MacDonald¹, Michael R. Heath¹, Simon P.R. Greenstreet², Douglas C. Speirs¹

¹Department of Mathematics & Statistics, University of Strathclyde, Glasgow, UK

²Marine Scotland Science, Marine Laboratory, Aberdeen, UK

***Correspondence:**

Alan MacDonald

Department of Mathematics & Statistics, University of Strathclyde, Glasgow, UK,
alan.macdonald@strath.ac.uk

Keywords: *Ammodytes marinus*¹, fish², larvae³, spawning⁴, hatching⁵, growth⁶, length⁷, 0-group

⁸

FIGURE CAPTIONS

Figure S1: Bootstrap estimates of the uncertainty in predicted hatch and spawning days. Each box-and-whisker plot shows the extremes (whiskers), quartiles (box) and median (horizontal line) of the estimated values for each year, as a consequence of imposing normally distributed variability in annual average growth rates.

TABLE CAPTIONS

Table S1: Bootstrap estimates of the uncertainty in predicted hatch and spawning days. Shown is the average interquartile range (days) of the estimated values for each hatch or spawning day between 2000 and 2009, as a consequence of imposing normally distributed variability in annual average growth rates.

TABLES

Table S1.

Hatch start interval	Hatch median interval	Hatch end interval	Spawning start interval	Spawning end interval
1	1.5	1.5	0.9	1.5

FIGURES

Figure S1.

