

Table S7 The upregulated proteins in both the 10% NaCl group compared with the control group and in the 20% NaCl group compared with the 10% NaCl group.

| Accession Number | Protein Name | Gene Name | Fold Change | |
|------------------|---|-------------------------|-------------|------------|
| | | | 10% NaCl ↑ | 20% NaCl ↑ |
| A0A077W129 | 30S ribosomal protein S13 | <i>rpsM</i> | 1.20 | 1.22 |
| A0A0D1FKQ9 | 30S ribosomal protein S9 | <i>rpsI</i> | 1.20 | 1.38 |
| A0A0D1H0K7 | 50S ribosomal protein L23 | <i>rplW</i> | 1.21 | 1.39 |
| A0A0B6XLK3 | Haloacid dehalogenase-like hydrolase | <i>ppaX</i> | 1.22 | 1.56 |
| A0A0D1H8Z3 | 3-oxoacyl-[acyl-carrier-protein] synthase 3 | <i>fabH</i> | 1.23 | 1.58 |
| A0A0D1HQY5 | Nucleoside diphosphate kinase | <i>ndk</i> | 1.24 | 1.48 |
| A0A0B6XQS8 | Valine-tRNA ligase | <i>valS</i> | 1.24 | 1.26 |
| A0A0D1I3J5 | Urease accessory protein UreE | <i>ureE</i> | 1.25 | 1.64 |
| A0A0D1HYX5 | 50S ribosomal protein L14 | <i>rplN</i> | 1.25 | 1.65 |
| A0A090N2E5 | Mannitol-specific phosphotransferase enzyme IIA component | <i>mtlF</i> | 1.28 | 1.62 |
| A0A0D1GWI4 | 50S ribosomal protein L4 | <i>rplD</i> | 1.28 | 1.35 |
| A0A0D1IW07 | Histidine-tRNA ligase | <i>hisS</i> | 1.29 | 1.61 |
| A0A0D1IHK1 | 50S ribosomal protein L20 | <i>rplT</i> | 1.33 | 1.50 |
| A0A090LYX3 | Inosine-uridine preferring nucleoside hydrolase | <i>SAU06011_2_40119</i> | 1.42 | 1.43 |
| A0A0D1JWV4 | 50S ribosomal protein L28 | <i>rpmB</i> | 1.55 | 1.65 |
| A0A0D1GP04 | Succinyl-CoA ligase [ADP-forming] subunit alpha | <i>SAJPND1_01177</i> | 1.49 | 2.10 |
| A0A0D6GUT4 | Membrane protein | <i>mmpL8</i> | 1.55 | 2.48 |
| A0A0D1H3E1 | Sced | <i>SAJPND1_02072</i> | 1.75 | 4.71 |
| A0A0B6XR64 | Urease subunit beta | <i>ureB</i> | 1.75 | 3.76 |
| A0A090LXA3 | Lipase/esterase LipA | <i>SAU06011_2_10612</i> | 1.80 | 2.06 |