

Berberine protects C17.2 neural stem cells from oxidative damage followed by inducing neuronal differentiation

Jia-Wen Shou ^{1,2}, Chun-Kai Cheung ², Jian Gao ⁴, Wei-Wei Shi ⁴ and Pang-Chui Shaw ^{1, 2, 3,*}

¹ School of Life Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong, China

² Li Dak Sum Yip Yio Chin R & D Centre for Chinese Medicine, The Chinese University of Hong Kong, Shatin, Hong Kong, China

³ State Key Laboratory of Research of Bioactivities and Clinical Applications of Medicinal Plants (CUHK) and Institute of Chinese Medicine, The Chinese University of Hong Kong, Shatin, Hong Kong, China

⁴ Shenzhen Health Development Research Center, Shenzhen, China

*Corresponding authors:

Wei-Wei Shi

Shenzhen Health Development Research Center, Shenzhen, China

Tel.: +86-755-83552132; Fax: +86-755-83552132.

Email address: shi1985@outlook.com

Pang-Chui Shaw

School of Life Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong, China,

Tel.: +852 39431363; Fax: +852 26037246.

Email address: pcshaw@cuhk.edu.hk

Supplementary Figure legends

FIGURE 1S| C17.2 cell viability after berberine (0.85, 1.69, 3.38, 6.75, 13.5, 27.0, 54.0 μM) treatment for 12 hours. The data represent the mean \pm SEM.

FIGURE 2S| C17.2 cell viability after berberine (0.85, 1.69, 3.38, 6.75, 13.5, 27.0, 54.0 μM) treatment for 24 hours. The data represent the mean \pm SEM. * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$.

Figure 1S




Figure 2S

