

Supplementary Figure 4

Prominence of IL6, IGF, TLR and bioenergetics pathway perturbation in lung tissues of scleroderma patients with pulmonary fibrosis

Ludivine Renaud¹, Willian A. da Silveira², Naoko Takamura¹, Gary Hardiman², Carol Feghali-Bostwick^{1*}


¹ Department of Medicine, Medical University of South Carolina, Charleston, SC, USA.

² School of Biological Sciences and Institute for Global Food Security, Queens University Belfast, Belfast BT9 5AG, UK.


*** Correspondence:**

Dr. Carol Feghali-Bostwick
feghalib@musc.edu

Gene expression for IGFBP2 (ENSG00000115457;5)



Gene expression for IGFL2 (ENSG00000204886;4)



Gtex is a comprehensive public resource to study tissue-specific gene expression and regulation. Samples were collected from 53 non-diseased tissue sites across nearly 1000 individuals, primarily for molecular assays including WGS, WES, and RNA-Seq.

Supplementary Figure 4: Tissue specific expression levels of IGFBP2 and IGFL2 in non-diseased organs. Figure generated by GTEx portal. Transformed fibroblasts, lung and skin (not sun-exposed and sun-exposed) are highlighted in red. TPM (Transcripts Per Kilobase Million)