


Tracking of an oral *Salmonella*-based vaccine for type 1 diabetes in non-obese diabetic mice

Jacques C. Mbongue¹, Ali Alhoshani², Jeffery Rawson¹, Pablo A. Garcia¹, Nelson Gonzalez¹, Kevin Ferreri¹, Fouad Kandeel¹, and Mohamed I. Hussein^{1,3,*}

¹Department of Translational Research & Cellular Therapeutics, Diabetes & Metabolism Research, Institute, Beckman Research Institute of City of Hope National Medical Center, Duarte, California 91010-3000, USA

²Department of Pharmacology and Toxicology, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia

³Faculty of Pharmacy, Zagazig University, Zagazig, Egypt



Supplementary Figure 1: A *Salmonella*-based vaccine does not impact B cell and polymorph nuclear macrophage numbers in lymphoid organs. NOD mice were injected with 2.5 μ g/mouse of anti-CD3 mAb for 5 days followed by 2 doses of oral combined PPI+TGF β +IL10 vaccine (n=5). Four weeks post-vaccination, B cells and PMN were analyzed in different organs by FACS analysis. **(A)** Quantification of the percentage of B220⁺GR-1⁻ (B cells) in total live cells of the spleen, PLN, MLN, and PP. **(B)** Quantification of the percentage of CD11b⁺GR-1⁺ (PMN) in total live cells of the spleen, PLN, MLN, and PP. The data shown are the average with SD from 2 independent experiments. Statistical analysis using two-way ANOVA shows significance between vaccine and vehicle treated groups (*p<0.01, ***P<0.001, ****p<0, 0001).