Supplementary information

Distribution of molecules related to neurotransmission in the nervous system of the mussel *Crenomytilus grayanus*

Elena Kotsyuba¹, Alexander Kalachev¹, Polina Kameneva¹& Vyacheslav Dyachuk^{1,2*}

Supplementary Figure 1. Whole structure of the main ganglia of the mussel *Crenomytilus grayanus*. (a) cerebropleural ganglion; (b) pedal ganglion; (c) visceral ganglion stained for 5-HT with DAPI. CPG, cerebropleural ganglion; PG, pedal ganglion; VG, visceral ganglion; CPPC, cerebral-pleural-pedal connectives; CPVC, cerebral-pleural-visceral connectives. Bar, 1mm.

Other Supplementary Information for this manuscript includes the following:

Supplementary Movie 1

Whole-mount immunostaining of the cerebropleural ganglion with antibody against 5-

HT.

Supplementary Movie 2

Whole-mount immunostaining of the cerebropleural ganglion with antibody against

Supplementary Movie 3

FMRFamide.

Whole-mount immunostaining of the pedal ganglion with antibody against 5-HT.

Supplementary Movie 4

¹ A.V. Zhirmunsky National Scientific Center of Marine Biology, Far Eastern Branch, Russian Academy of Sciences, Vladivostok 690041, Russia

²Department of Nanophotonics and Metamaterials, IFMO University, St. Petersburg, Russia

^{*}Corresponding author: slavad83@gmail.com

Whole-mount immunostaining of the pedal ganglion with antibody against FMRFamide.

Supplementary Movie 5

Whole-mount immunostaining of the visceral ganglion with antibody against FMRFamide.