


*Supplementary Material***Identification of Genetic Variants via Bacterial Respiration Gas Analysis**

**Naoki Koga, Takuro Hosomi, Martijn Zwama, Chaiyanut Jirayupat, Takeshi Yanagida, Kunihiko Nishino\*, Seiji Yamasaki\***

**\* Correspondence:** Kunihiko Nishino. [nishino@sanken.osaka-u.ac.jp](mailto:nishino@sanken.osaka-u.ac.jp)  
Seiji Yamasaki. [seiji37@sanken.osaka-u.ac.jp](mailto:seiji37@sanken.osaka-u.ac.jp)

## Supplementary Figures



**Supplementary Figure 1. Repeats of the GC-MS chromatograms of the gas released from different *Escherichia coli* strains.** Chromatograms of the gas present in the absorbent for all four strains MC4100, MG1655, MC4100 $\Delta$ tnaAB, and MG1655 $\Delta$ tnaAB. The detected organic species were identified from their mass/charge ratio.

**Supplementary Table 1**

	MC4100	MG1655	MC4100 <i>ΔtnaAB</i>	MG1655 <i>ΔtnaAB</i>
Amount of adsorbed indole (nmol)	1.5	1.4	0	0

Adsorbed amount of indole calculated by GC-MS analyses using a calibration standard.