Supplementary table S3. Hematology of preterm pigs treated with rhIGF-1/BP-3 or vehicle*

| Parameter | rhIGF-1/BP-3 | Controls |
|--|-------------------|-------------------|
| Number of animals | 20 | 21 |
| Total leukocytes, 10 ⁹ /L | 2.4 ±0.2* | 1.8 ±0.1 |
| Total erythrocytes, 10 ⁹ /L | 3.8 ± 0.1 | 3.6 ± 0.1 |
| Thrombocytes, 10 ⁹ /L | 179.2 ±11.3 | 172.4 ±13.4 |
| Neutrophils, 10 ⁹ /L | 1.2 ±0.1* | 0.8 ± 0.1 |
| Lymphocytes, 10 ⁹ /L | 1.1 ±0.1 | 0.9 ± 0.1 |
| Monocytes, 10 ⁹ /L | 0.04 ± 0.01 | 0.03 ± 0.003 |
| Eosinophils, 10 ⁹ /L | 0.04 ± 0.01 | 0.04 ± 0.01 |
| Basophiles, 10 ⁹ /L | $5e-04 \pm 0.001$ | 0.001 ± 0.001 |
| LUC, 10 ⁹ /L | 0.01 ± 0.002 | 0.01 ± 0.002 |
| Hemoglobin, mmol, L | 5.3 ±0.1 | 5.1 ± 0.1 |
| Hematocrit, L/L | 0.3 ±0.01 | 0.3 ± 0.05 |
| MCV, ft | 71.0 ± 0.9 | 70.0 ± 0.6 |
| MCHC, mmol/L | 19.8 ± 0.8 | 19.9 ± 0.1 |
| Mean platelet volume, ft | 11.6 ± 0.5 | 11.0 ± 0.2 |
| Mean platelet count, g/L | 227.9 ±2.3 | 226.0 ± 2.4 |
| Neutrophils, % | 49.0 ± 2.3 | 46.3 ± 2.7 |
| Lymphocytes, % | 46.8 ± 2.3 | 49.4 ± 2.6 |
| Monocytes, % | 1.8 ± 0.2 | 1.9 ± 0.2 |
| Eosinophils, % | 1.7 ± 0.2 | 1.8 ±0.3 |
| Basophils% | 0.08 ± 0.01 | 0.15 ± 0.07 |
| LUC, % | 0.5 ±0.1 | 0.4 ±0.1 |

^{*}Values are mean \pm SEM. MCV, mean corpuscular volume. There were no statistically significant differences in the variables (p > 0.05), except that rhIGF-1/BP-3 treatment increased leukocyte and neutrophil counts (p < 0.05). MCHC, mean corpuscular hemoglobin concentration LUC, large unstained cells.