

Table S4: Strains and plasmids used in this study.

Strain or plasmid	Relevant characteristic(s) ^a	Reference or source
<i>P. aeruginosa</i>		
Wild type	PAO1 wild type	laboratory collection
$\Delta czcR\Delta czcS$	PAO1 $\Delta czcR\Delta czcS$	(1)
$\Delta copR\Delta copS$	PAO1 $\Delta copR\Delta copS$	This study
$\Delta PA2807$	PAO1 $\Delta PA2807$	This study
<i>E. coli</i>		
DH5 α	recA1, endA1, hsdR17, deoR, thi-1, supE44, gyrA96, relA1, Δ (lacZYA-argF), U169(φ 80dlacZ Δ M15)	(2)
BL21(DE3)	E. coli str. B F- ompT gal dcm lon hsdSB(rB-mB-) λ (DE3 [lacI lacUV5-T7p07 ind1 sam7 nin5]) [malB+]K-12(λ S)	(3)
Plasmids		
pME3087	Suicide plasmid, CoIE1 replicon; Tc ^r	(4)
pMMB66EH	Expressing vector carrying an IPTG-inducible promoter; Ap ^r , Cb ^r	(5)
pMMB66EH- <i>PA2807</i>	pMMB66EH derivative, carrying the <i>PA2807</i> gene; Ap ^r , Cb ^r	This study
pME6001	pME6000 derivative plasmid; Gm ^r	(6)
pME6001- <i>PA2807</i>	pME6001 carrying the <i>PA2807</i> gene and its promoter; Gm ^r	This study
pBBR1- <i>gfp</i>	Transcriptional <i>gfp</i> fusion cloning vector; Ap ^r , Cb ^r	(7)
<i>czcC:gfp</i>	pBBR1 derivative, carrying the <i>czcC</i> promoter; Ap ^r , Cb ^r	(8)
<i>czcD:gfp</i>	pBBR1 derivative, carrying the <i>czcD</i> promoter; Ap ^r , Cb ^r	This study
pGex-2T- <i>zur</i>	GST-fusion expression plasmid, carrying the <i>zur</i> gene; Ap ^r	This study

^a Antibiotic resistance is indicated by r: Ap ampicillin; Tc tetracycline; Cb carbenicillin; Gm gentamycin

References:

1. Caille, O., Rossier, C. and Perron, K. (2007) A copper-activated two-component system interacts with zinc and imipenem resistance in *Pseudomonas aeruginosa*. *J Bacteriol*, **189**, 4561-4568.
2. Sambrook, J., and D. W. Russell. . (2001.) *Molecular cloning: a laboratory manual* 3rd ed.Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
3. Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Methods in enzymology*, **185**, 60-89.
4. Voisard, C., Bull, C.T., Keel, C., Laville, J., Maurhofer, M., Schnider, U., Défago, G. and Haas, D. (1994), *Molecular Ecology of Rhizosphere Microorganisms*, pp. 67-89.

5. Furste, J.P., Pansegrau, W., Frank, R., Blocker, H., Scholz, P., Bagdasarian, M. and Lanka, E. (1986) Molecular cloning of the plasmid RP4 primase region in a multi-host-range tacP expression vector. *Gene*, **48**, 119-131.
6. Blumer, C., Heeb, S., Pessi, G. and Haas, D. (1999) Global GacA-steered control of cyanide and exoprotease production in *Pseudomonas fluorescens* involves specific ribosome binding sites. *Proc Natl Acad Sci U S A*, **96**, 14073-14078.
7. Ouahrani-Bettache, S., Porte, F., Teyssier, J., Liautard, J.P. and Kohler, S. (1999) pBBR1-GFP: a broad-host-range vector for prokaryotic promoter studies. *Biotechniques*, **26**, 620-622.
8. Ducret, V., Gonzalez, M.R., Leoni, S., Valentini, M. and Perron, K. (2020) The CzcCBA Efflux System Requires the CadA P-Type ATPase for Timely Expression Upon Zinc Excess in *Pseudomonas aeruginosa*. *Front Microbiol*, **11**, 911.