

Supplementary materials

Table S1. Information criteria and entropy for latent class solutions 2 to 5

| Number of Classes | AIC | BIC | Entropy | n smallest class |
|----------------------|----------|----------|---------|---------------------|
| 2 | 2001.707 | 2046.307 | 0.948 | 27 |
| 3 | 1945.719 | 2007.044 | 0.92 | 17 |
| 4 | 1901.896 | 1979.946 | 0.934 | 15 |
| 5 | 1867.587 | 1962.361 | 0.954 | 7 |

Note: AIC: Akaike Information Criteria BIC: Bayesian Information Criteria

Table S2. Properties of the latent trajectory classes: sociodemographic factors, depression, and prior psychiatric diagnosis

| Sociodemographics | | | | | | |
|---------------------------------------|----------------|-------|----------------------|-------------------|--------------------|-----------------|
| | Sex (%male) | age | Employment (%yes) | partner (%yes) | married (% yes) | child (%yes) |
| Class 1 | 45.1% | 38.93 | 59.2% | 55.9% | 41.5% | 47.6% |
| Class 2 | 33.9% | 35.75 | 51.8% | 26.0% | 1.4% | 23.4% |
| Class 3 | 52.6% | 35.86 | 60.8% | 23.7% | 8.1% | 11.1% |
| Depression (BDI) score | | | | | | |
| | t1 | t2 | t3 | t4 | t5 | |
| Class 1 | 14.86 | 7.83 | 7.55 | 6.31 | 4.00 | |
| Class 2 | 28.66 | 26.39 | 20.46 | 25.69 | 19.46 | |
| Class 3 | 21.31 | 20.57 | 17.00 | 22.69 | 12.45 | |
| Prior diagnosis (ICD-10 codes) | | | | | | |
| | F1 | F3 | F4 | F6 | others | |
| Class 1 | 25.9% | 59.6% | 48.8% | 8.1% | 6.6% | |
| Class 2 | 22.8% | 70.1% | 34.2% | 42.9% | 7.6% | |
| Class 3 | 25.4% | 68.8% | 37.9% | 23.0% | 1.7% | |

(ICD-10) codes: F1*, substance abuse disorder; F3*, affective disorder; F4*, neurotic and acute stress reaction; F6*, personality disorder.